


MULTI-GRID CONTINUATION AND SPURIOUS SOLUTIONS FOR  
NONLINEAR BOUNDARY VAL. (U) ARIZONA UNIV TUCSON DEPT OF  
MATHEMATICS H MITTELMANN JUL 86 TR-90 AFOSR-TR-86-0888  
AFOSR-84-0315 F/G 12/1

ML

**UNCLASSIFIED**

F/G 12/1



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AFOSK-TR- 86-0888

Multi-grid Continuation and  
Spurious Solutions for Nonlinear  
Boundary Value Problems

by

H. D. Mittelman

Technical Report No. 90  
January 1986

Arizona State University  
Tempe, AZ 85287

Approved for public release;  
distribution unlimited.

# DEPARTMENT OF MATHEMATICS

DTIC  
ELECTE  
S OCT 08 1986 D



AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)  
NOTICE OF TRANSMITTAL TO DTIC  
This technical report has been reviewed and is  
approved for public release (AFOSR 19-1-22)  
Distribution is unlimited.  
MATTHEW J. KERPEN  
Texas Tech. Tech. Information Division

ARIZONA STATE UNIVERSITY

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

AD-A173 081

DTIC FILE COPY

Multi-grid Continuation and  
Spurious Solutions for Nonlinear  
Boundary Value Problems

by

H. D. Mittelmann

Technical Report No. 90  
January 1986

Arizona State University  
Tempe, AZ 85287

DTIC  
ELECTE  
S OCT 08 1986 D  
D

Arizona State University is an affirmative action/equal employment opportunity institution and does not discriminate on the basis of sex, race, ethnic origin, creed, color, age, handicapped or veteran status in its program or employment.

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

AD-A173081

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY ---			3. DISTRIBUTION AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A			5. MONITORING ORGANIZATION REPORT NUMBER <b>AFOSR-TR- 86-0888</b>		
4. PERFORMING ORGANIZATION REPORT NUMBER			7a. NAME OF MONITORING ORGANIZATION <b>AFOSR</b>		
6a. NAME OF PERFORMING ORGANIZATION <b>Arizona State University</b>		6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State and ZIP Code) <b>Building 410 Bolling AFB DC 20332-6448</b>		
6c. ADDRESS (City, State and ZIP Code) <b>Temple AZ 85287</b>		8a. NAME OF FUNDING/SPONSORING ORGANIZATION <b>AFOSR</b>			
8b. OFFICE SYMBOL (If applicable) <b>NM</b>		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER <b>AFOSR 84-0315</b>			
8c. ADDRESS (City, State and ZIP Code) <b>Building 410 Bolling AFB DC 20332-6448</b>		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO. <b>61102F</b>	PROJECT NO. <b>2304</b>	TASK NO. <b>A3</b>	WORK UNIT NO.
11. TITLE (Include Security Classification) <del>CONTINUATION AND</del> <del>MULTI GRID METHODS FOR BIFURCATION PROBLEMS</del> <i>See title page</i>					
12. PERSONAL AUTHOR(S) <b>Dr Hans Mittelmann</b>					
13a. TYPE OF REPORT <b>ANNUAL</b>		13b. TIME COVERED FROM <b>10/01/85</b> TO <b>7/01/86</b>		14. DATE OF REPORT (Yr., Mo., Day) <b>Jan. 1986</b>	
15. PAGE COUNT <b>26</b>		16. SUPPLEMENTARY NOTATION <b>July per phone call</b>			
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The Investigator has employed multi-grid methods to solve an important class of boundary value problems in two space dimensions that arises in VLSI device simulation. Methods for the numerical solution of bifurcation problems are also under development. Six papers have been produced so far under this effort.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <b>UNCLASSIFIED/UNLIMITED</b> <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		
22a. NAME OF RESPONSIBLE INDIVIDUAL <b>JOHN P. THOMAS, CAPT, USAF</b>		22b. TELEPHONE NUMBER (Include Area Code) <b>202/767-5026</b>		22c. OFFICE SYMBOL <b>NM</b>	

DD FORM 1473, 83 APR

86

EDITION OF 1 JAN 79 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

( 4 )

Multi-grid Continuation and Spurious Solutions  
for Nonlinear Boundary Value Problems

H. D. Mittelmann

Department of Mathematics  
Arizona State University  
Tempe, Arizona 85287

Abstract Recently the author together with R. Bank has developed, implemented and successfully applied a continuation technique for the numerical solution of parameter-dependent nonlinear elliptic boundary value problems. The method was integrated into an existing multi-grid package based on an adaptive finite element discretization. <sup>This document</sup> We present the continuation method and prove an important theoretical result for the corrector iteration. For the Bratu problem <sup>the model problem</sup>  $- \Delta u = \lambda e^u$  on the square with homogeneous Dirichlet conditions <sup>it is</sup> we show how spurious solutions may be encountered while computing relevant solutions, how the program handles those and how it allows to detect them.

This work was supported by the Air Force Office of Scientific Research under Grant AFOSR-84-0315.

<input checked="checked" type="checkbox"/>	Codes
<input type="checkbox"/>	Major
<input type="checkbox"/>	Special
A-1	

## 1. Introduction

In the following we consider the parameter-dependent nonlinear problem

$$(1.1) \quad G(u, \alpha) = 0,$$

where  $G : X^m \times \mathbb{R}^p \rightarrow X^m$ ,  $X$  a suitable function space and  $\alpha$  a vector of real parameters. (1.1) will typically represent a parameter-dependent nonlinear elliptic system of dimension  $m$  and order two. Important examples are the VLSI device simulation equations

$$g_1(u, v, w) \equiv -\Delta u + e^{u-v} - e^{w-u} - k_1 = 0,$$

$$(1.2) \quad g_2(u, v, w) \equiv \nabla \cdot \mu_n e^{u-v} \nabla v + k_2 = 0,$$

$$g_3(u, v, w) \equiv -\nabla \cdot \mu_p e^{w-u} \nabla w + k_2 = 0.$$

In this case the parameters  $\alpha$  enter the boundary conditions if, for example, current-voltage characteristics are to be determined (cf.[3]).

The solution manifold  $G^{-1}(0)$  of (1.1) may have a very complex structure. In practical applications it is frequently desirable and sufficient to compute certain cross-sections of it, i.e. all parameters  $\alpha$  are kept fixed at certain values and for simplicity suppressed except one which we will call  $\lambda$ . A simple graphical representation of this cross-section may be obtained if a functional of the solution is depicted versus  $\lambda$ . It should be noted, however, that usually in these applications such functionals are in fact important pieces of information, such as the currents for (1.2).

We assume that a vector

$$(1.3) \quad R(u(\lambda)), \quad R : X^m \rightarrow \mathbb{R}^k$$

of differentiable functionals is given and that most of the important information about the problem can be obtained from the  $k \times p$  diagrams

$$(1.4) \quad R_i = R_i(u(\alpha_j)), \quad i = 1, \dots, k, \quad j = 1, \dots, p$$

and the corresponding solutions.

The solutions  $u(\lambda)$ ,  $\lambda = \alpha_j$ , in (1.4) and from them the curves  $R_i(u(\lambda))$  usually have to be computed in a continuation fashion. Starting from a known solution  $u(\lambda_0)$  another solution is computed by utilizing a predictor-corrector scheme. Frequently an initial solution to start this continuation procedure is not known and has itself to be determined from an approximate solution. Along the solution curves singular points have to be expected and our goal is to provide methods to locate these points, to overcome turning points and to switch branches at bifurcation points.

In the following section we will outline a continuation algorithm that has been successfully used for the scalar case ( $m = 1$  in (1.1)). It provides an efficient and robust method and has been implemented for a general class of second order elliptic boundary value problems discretized by finite elements in [2]. The continuation method is applied on a coarse triangulation of the given domain and fine grid approximations at arbitrarily specified points are obtained by multi-grid.



## 2. The Continuation Method

In the following we assume that a choice of the parameter  $\lambda = \alpha_j$  in (1.1), (1.3) and also of the functional  $R_1$  in (1.4) has been made and denote (1.1) by

$$(2.1) \quad G(u, \lambda) = 0, \quad G : X^m \times \mathbb{R} \rightarrow X^m,$$

and  $R_1$  by  $r$ .

The parametrization of the solution  $u$  of (2.1) by  $\lambda$  and thus of  $r$  by  $\lambda$  need not be possible in general. Let  $s$  denote the arclength along a solution curve  $(u, \lambda) = (u(s), \lambda(s))$  of (2.1). Then under sufficient smoothness and regularity assumptions this latter dependence is differentiable and thus  $(G_u^0 = G_u(u_0, \lambda_0)$  etc.).

$$(2.2a) \quad G_u^0 \dot{u}_0 + G_\lambda^0 \dot{\lambda}_0 = 0,$$

$$(2.2b) \quad \|\dot{u}_0\|^2 + \dot{\lambda}_0^2 = 1$$

in a solution  $(u_0, \lambda_0) = (u(s_0), \lambda(s_0))$  of (2.1). Here, subscripts stand for partial derivatives and a dot for differentiation with respect to  $s$ .

In singular points  $(u_0, \lambda_0)$  of (2.1) the linear operator  $G_u^0$  in (2.2a) is singular and, for example, a solution by Newton's method breaks down. This singularity may be removed or reduced by augmenting (2.1) by a normalizing equation, a well-known example of which is

$$(2.3) \quad N_\sigma(u, \lambda, \sigma) \equiv \theta \dot{u}_0(u - u_0) + (2 - \theta) \dot{\lambda}_0(\lambda - \lambda_0) - \delta \sigma = 0$$

which approximates (2.2b).  $\sigma$  is called the pseudo-arclength parameter.  $\theta$  satisfying  $0 \leq \theta \leq 2$  may be used as a weighting parameter.

Instead of (2.3) we propose to use

$$(2.4) \quad N_r(u, \lambda, \sigma) \equiv \theta \dot{r}_0 (r - r_0) + (2 - \theta) \dot{\lambda}_0 (\lambda - \lambda_0) - \delta \sigma = 0,$$

where  $r_0 = r(u_0)$ ,  $\dot{r}_0 = \left( \frac{d}{d\sigma} r(u) \right)_{\sigma = \sigma_0}$  and  $r$  denotes the above functional. From (2.4) we see that the three values  $\theta = 0, 1, 2$  are special in the sense that for  $\theta = 0(2)$  the augmenting equation characterizes points with a fixed  $\lambda(r)$  - value, while for  $\theta = 1$  this point lies on a hyperplane orthogonal to the solution arc in  $(r_0, \lambda_0)$ .

This suggests to use target values in the two (physically relevant) variables  $r$  and  $\lambda$ . Such a strategy seems appropriate as long as starting from a point on the solution curve another point corresponding to a suitably chosen target value can be computed in a small number of (corrector) iterations. This was achieved for the scalar case of (2.1) and the choice

$$(2.5) \quad r(u) = \|u\|.$$

The following class of nonlinear boundary value problems in the plane

$$(2.6) \quad \begin{aligned} -\nabla \cdot a(x, y, u, \nabla u, \lambda) + f(x, y, u, \nabla u, \lambda) &= 0 && \text{in } \Omega, \\ u &= g_1(x, y, \lambda) && \text{on } \partial\Omega_1, \\ a \cdot n &= g_2(x, y, u, \lambda) && \text{on } \partial\Omega_2 = \partial\Omega \setminus \partial\Omega_1, \end{aligned}$$

was considered. Here,  $\Omega$  is a connected domain in  $R^2$  with boundary  $\partial\Omega$ ,  $n$  is the

unit normal vector on  $\partial\Omega$ ,  $a = (a_1, a_2)^T$  and  $a_1, a_2, f, g_1, g_2$  are given scalar functions.

In [2] the program PLTMG, an implementation of an adaptive finite element multi-grid method combined with the continuation technique presented below is described in detail. In addition to some theoretical results on the continuation method [12, 13] contain numerical results for several problems from the applications including problems with turning-points, symmetry-breaking and other bifurcation points as well as parameter-switching for cases depending on several ( $p > 1$  in (1.1)) parameters. [13] also contains a comparison with the classical pseudo-arclength continuation method. Other problems to which the program has been applied successfully are Steklov eigenvalue problems for which the Dirichlet condition in (2.6) is parameter-dependent.

The predictor step which is crucial for the performance of the method determines predicted values of the form

$$(2.7) \quad \begin{pmatrix} u_p \\ \lambda_p \end{pmatrix} = \begin{pmatrix} (1+\gamma)u_0 + \beta \dot{u}_0 \\ \lambda_0 + \alpha \dot{\lambda}_0 \end{pmatrix}$$

such that  $\alpha, \beta, \gamma$  satisfy the three equations

$$(2.8) \quad \begin{aligned} N(u_p, \lambda_p, \sigma) &= 0, \\ (u_0, G(u_p, \lambda_p)) &= 0, \\ (\dot{u}_0, G(u_p, \lambda_p)) &= 0 \end{aligned}$$

in a least-squares sense. Here it is assumed that  $X$  is a Hilbert space and  $(\cdot, \cdot)$  denotes its inner product.

In the corrector iteration the augmented system

$$(2.9) \quad F(y) = \begin{pmatrix} G(u, \lambda) \\ N(u, \lambda, \sigma) \end{pmatrix} = 0, \quad y^T = (u^T, \lambda)$$

is solved by Newton's method starting in  $(u_p, \lambda_p)$ . The restriction of the problem (2.1) to the  $(r, \lambda)$ -plane in case  $N = N_r$  is used in (2.8) introduces an additional singularity in exceptional points.

If in a solution  $(u_0, \lambda_0)$  simultaneously  $\dot{r}_0 = 0$  and  $\dot{\lambda}_0 = 0$  then the normalizing condition (2.4) is not well-defined. This situation occurs in symmetry-breaking pitchfork bifurcation points, see [11]. We propose to switch temporarily to the augmenting equation (2.3) which, as explained in [12] may be viewed as asymptotically equivalent to a version of the generalized inverse iteration ([10]) exploiting symmetries in such bifurcation points. The precise choice of  $N$  is given by (cf. [13]).

$$|\dot{r}_0| < \epsilon \|\dot{u}_0\| : N = N_\sigma, \theta \neq 0,$$

$$(2.10) \quad |\dot{r}_0| > (1-\epsilon) \|\dot{u}_0\| : N = N_r, \theta \neq 0, \beta = 0,$$

$$\epsilon \|\dot{u}_0\| \leq \dot{r}_0 \leq (1-\epsilon) \|\dot{u}_0\| : N = N_r, 0 \leq \theta \leq 2.$$

### 3. The Corrector Iteration

A point  $(u_p, \lambda_p)$  computed by the predictor of the previous section is used as a starting guess for an iterative solution of the augmented system (2.9). This is done by Newton's method and thus the Jacobian

$$(3.1) \quad F_y = \begin{pmatrix} G_u & G_\lambda \\ N_u & N_\lambda \end{pmatrix}$$

has to be evaluated and inverted.

While (2.10) prescribes which  $N$  to choose it does not completely define the augmenting equation. The weighting parameter  $\theta$  still has to be picked and we assume that  $r$  respectively  $\lambda$  is not used as parameter to continue past a turning point in  $r$  respectively  $\lambda$ . This yields  $\theta \neq 0$  if  $\dot{\lambda}_0 = 0$  and  $\theta \neq 2$  if  $\dot{r}_0 = 0$ . Under these assumptions it was shown in Proposition 4.2 of [13] that  $F_y$  is regular except in bifurcation points.

When computing in a neighborhood of a bifurcation point the growth of  $\|F_y^{-1}\|$  is crucial for the convergence of Newton or other iterative methods used as corrector, cf. [7]. We give here the detailed proof of Proposition 4.3 in [13]. Let locally near the simple bifurcation point  $y_0 = (u_0, \lambda_0)$  solutions be parametrized by  $s$  such that without restriction of generality  $y_0 = (u(0), \lambda(0))$ . We assume in the following that  $\dot{y}_0 = (\dot{u}(0), \dot{\lambda}(0))$  is a nondegenerate ray in the sense of [8].

Theorem 3.1 If the normalizing condition  $N$  is chosen according to (2.10) then on any of the branches passing through the simple bifurcation point  $y_0 = (u(0), \lambda(0))$  of (2.1) for  $\delta > 0$  sufficiently small

$$(3.2) \quad \|F_y^{-1}(y(s))\| = O(|s|^{-1}), \quad 0 < |s| < \delta.$$

Proof. As mentioned in [13] we have to generalize the proof of Theorem 1 in [8]. This proof covers the case  $N = N_\sigma$  so that it remains to consider  $N = N_r$ . We note that the row-vector  $a_0 = (N_u, N_\lambda)$  in (3.1) is not constant in this case, in fact

$$(3.3) \quad a_0(s) = \left( \frac{\theta \dot{r}_0 u(s)}{\|u(s)\|}, (2-\theta)\dot{\lambda}_0 \right).$$

Let us further denote  $H(y) = G(u, \lambda)$  and thus

$$H' = (G_u, G_\lambda) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n, \quad F_y = \begin{pmatrix} H' \\ a_0 \end{pmatrix}.$$

As in [8] we introduce the one-dimensional space  $X_1 \subset \mathbb{R}^{n+1}$  spanned by  $\dot{y}_0 = (\dot{u}_0, \dot{\lambda}_0)^T$  on one of the branches, the one-dimensional subspace  $X_2 \subset \mathbb{R}^{n+1}$  such that the kernel  $K(H'(y_0))$  satisfies  $K = X_1 \oplus X_2$ , and the  $(n-1)$ -dimensional subspace  $X_3$  such that  $\mathbb{R}^{n+1} = X_1 \oplus X_2 \oplus X_3$ . Similarly let  $Y_1 = R$ ,  $Y_3 = R(H'(y_0))$ , where  $R(A)$  denotes the range of the linear operator  $A$ , and let  $Y_2$  be the one-dimensional subspace of  $\mathbb{R}^n$  such that  $\mathbb{R}^n = Y_2 \oplus Y_3$ . We define the orthogonal projectors  $Q : \mathbb{R}^n \rightarrow Y_3$  and  $I - Q : \mathbb{R}^n \rightarrow Y_2$ . For  $0 < |s| < \delta$  we define the linear operator  $B(s) : X_1 \times X_2 \times X_3 \rightarrow Y_1 \times Y_2 \times Y_3$  by

$$B(s)(x_1, x_2, x_3) = (a_0(s)(x_1 + x_2 + x_3), (I - Q)H'(y(s))(x_1 + x_2 + x_3), QH'(y(s))(x_1 + x_2 + x_3)).$$

$B(s)$  is equivalent to  $F_y(y(s))$  in the sense that

$$F_y = C B D$$

where  $C$  and  $D$  are regular matrices independent of  $s$ .  $B(s)$  can be decomposed in a natural way into nine suboperators  $B_{ij}(s) : X_i \rightarrow Y_j$ ,  $1 \leq i, j \leq 3$ . From the above definitions it follows that  $B_{21}(0)$ ,  $B_{22}(0)$ ,  $B_{31}(0)$ ,  $B_{32}(0)$  and  $B_{23}(0)$  vanish identically. From (2.10) we conclude the crucial fact

$$a_0(0)\dot{y}_0 = \theta \dot{r}_0^2 + (2-\theta)\dot{\lambda}_0^2 \neq 0$$

This and the definition of  $X_3$ ,  $Y_3$  imply that  $B_{11}(0)$  and  $B_{33}(0)$  define bijections, while as in [8] it follows that  $\dot{B}_{21}(0) = 0$  and that  $\dot{B}_{22}(0)$  is a bijection between  $X_2$  and  $Y_2$ . We can thus expand  $B(s)$  in a neighborhood of  $s = 0$  as

$$B(s) = \begin{pmatrix} B_{11}(0) & B_{12}(0) & B_{13}(0) \\ s^2 \tilde{B}_{21}(0) & s \dot{B}_{22}(0) & s \tilde{B}_{23}(0) \\ s \tilde{B}_{31}(0) & s B_{32}(0) & B_{33}(0) \end{pmatrix} + \text{h.o.t.}$$

where  $B_{ij}(0)$ ,  $\dot{B}_{ij}(0)$  and  $\tilde{B}_{ij}(0)$  are linear operators from  $X_i$  into  $Y_j$  which are uniformly bounded with respect to  $s$ . Thus  $\det B(s) = O(|s|)$  and hence  $B^{-1}(s)$  can be written as

$$B^{-1}(s) = \begin{pmatrix} E_{11}(s) & s^{-1}E_{12}(s) & E_{13}(s) \\ sE_{21}(s) & s^{-1}E_{22}(s) & E_{23}(s) \\ sE_{31}(s) & E_{32}(s) & E_{33}(s) \end{pmatrix}$$

where similarly the  $E_{ij}$  are uniformly bounded and  $E_{22}(0) = (\dot{B}_{22}(0))^{-1}$  is regular.

#### 4. Spurious Solutions

It is well-known that discretized problems possess frequently many more solutions than the continuous problems. All discrete solutions that do not correspond to solutions of the continuous problem and disappear for  $h \rightarrow 0$ ,  $h$  the discretization parameter, are called spurious or numerically irrelevant solutions. If we consider the simple model equation

$$(4.1) \quad -\Delta u = \lambda f(u)$$

then there are essentially two types of those solutions. One type bifurcating from  $\lambda = \pm \infty$  in case  $f(u)$  has zeros and the other bifurcating from  $\|u\| = \infty$  if the growth of  $f$  is sufficiently strong. It might be argued that not too much effort should be spent on investigating those spurious solutions if the main interest is that of approximately solving the continuous problem. There are, however, at least two counter arguments.

First, in a multi-grid context where one hopes to be able to use rather crude coarse grids on which the continuation will be done, one has to be aware of the spurious solutions since for 'large'  $h$  they are present for 'small' values of  $\|u\|$  respectively  $\lambda$  and thus may be more easily encountered. Second, it has been shown recently ([9]) that for the second type of spurious solutions some of these may bifurcate from a relevant branch. Thus they definitely will be encountered and it becomes necessary to distinguish between spurious and relevant solutions.

If we specialize (4.1) to

$$(4.2) \quad -\Delta u = \lambda e^u \quad \text{in } \Omega = [0,1]^n, \quad u = 0 \text{ on } \partial\Omega$$



then the proof in [9] covers the case  $n = 1$  and a uniform grid of an even number of points. The midpoint of  $[0,1]$  where the relevant solution attains its maximum is then not a grid point. The discrete solution attains its maximum in the two neighboring points and from this branch bifurcate two branches attaining their maximum in either of these points.

We had observed these solutions and their bifurcation previously when numerically solving (4.2) for  $n = 2$  by discretizing with the standard difference method on a corresponding grid. We also have applied a multi-grid method to compute spurious solutions ([14]). Since our goal is not to contribute to the knowledge about the spurious solutions but rather to provide a general purpose and if possible foolproof program to solve a general class of problems, it is natural to investigate the above situation with the program PLTMG.

Our expectation is that the implemented continuation method allows us to follow the spurious solution branches because these are genuine solutions of the discrete problems. On the other hand the multi-grid process using several discretizations simultaneously could help to identify spurious solutions. But additionally all the other features offered by the program as the adaptive refinement and the error estimators will have their impact on the solution process respectively may give other hints about which solutions are relevant.

Since the program uses a finite element discretization with piecewise linear elements we try to come close to the above discretization. The unit square is subdivided into nine subsquares of length  $h = 1/3$  and then all these squares are divided into two triangles by the SW - NE diagonal. This way the Laplace operator will be discretized in the same way as by the five point star difference method. The right-hand side, however, in the finite element context discretized by a quadrature formula will be different from the pointwise

discretization of the difference method. This will introduce a bias in the SW-NE direction and we do not expect the relevant solution to attain its maximum in the four interior nodes.

While the numerical results presented in the following, of course, make no statements about the mathematical properties of the problem they provide some valuable insight into the practical use of the program and the numerical solution in the presence of spurious solutions.

The branch starting at  $\lambda = 0$ ,  $r = \|u\| = 0$  for the above discretization with four nodes attains its maximum, as expected, in the points  $(1/3, 1/3)$ ,  $(2/3, 2/3)$  and has two equal but smaller values at  $(1/3, 2/3)$ ,  $(2/3, 1/3)$ . When following this curve beyond the limit point another sign-change of  $\det(G_u)$  is encountered. Even a tracing of the neighborhood of this point with small continuation steps shows an abrupt change of the sign. When asked to locate this point the program ends up on a different branch and even a subsequent branch switching does not lead back to the relevant branch. This suggests that instead of bifurcation a perturbed bifurcation seems to be present. This is not so relevant here, since the fact remains that a sign-change seems to indicate bifurcation and naturally a spurious branch is found. Table 4.1 shows the continuation to and the location of the 'bifurcation' point. In the tables NI denotes the number of corrector iterations and  $\delta$  is a quantity that changes sign at simple bifurcation points (cf. [2]).

In the following we present multi-grid results for four different cases. First we have followed the relevant branch up to  $r = 4.5$  respectively  $r = 5$  and computed a 3-grid solution for fixed norm with uniform and adaptive refinement. Table 4.2 contains only the refinement results.

NI	$\lambda$	$\ u\ $	$\dot{\lambda}$	$\dot{r}$	$\det(G_u)$	$\delta$
Continue to $\lambda = .1$ then to $r = 1, r = 3, r = 3.44, r = 3.46$ .						
1	0.000E+00	0.000E+00	0.995E+00	0.000E+00	0.492E 5	0.200E+01
1	0.100E+01	0.116E+00	0.992E+00	0.128E+00	0.276E 5	0.180E+01
1	0.390E+01	0.100E+01	0.244E+00	0.970E+00	0.190E 3	0.129E+00
2	0.156E+01	0.300E+01	-0.756E+00	0.653E+00	-0.623E 3	0.592E+00
2	0.111E+01	0.344E+01	-0.669E+00	0.741E+00	-0.185E 2	0.138E-01
1	0.109E+01	0.346E+01	-0.664E+00	0.745E+00	0.176E 2	-0.130E-01
Determinant changes sign; find bifurcation point by secant method on $\delta$ .						
1	0.110E+01	0.345E+01	-0.121E-01	0.130E-01	-0.113E -1	0.809E-05
1	0.110E+01	0.345E+01	0.192E+00	-0.215E+00	0.123E 0	-0.883E-04
1	0.110E+01	0.345E+01	0.497E-06	-0.591E-03	0.482E -2	-0.369E-09

Table 4.1 Continuation along relevant branch, detection of spurious branch.

A posteriori error estimates based on solving local problems with higher-order elements [5] are evaluated by the program and in Table 4.4 we give the number of estimated correct digits in three different norms.

Following the irrelevant branch that attains its maximum at  $(1/3, 1/3)$  to  $r = 5.5$  and again refining uniformly and adaptively yielded the results in Table 4.3. We see that for the adaptive refinement it seems impossible to conclude from the error estimations alone that the solution is irrelevant, while for the uniform refinement the error was estimated as bigger than the solution resulting in a negative number of estimated correct digits. In this latter case also the solution on the finest grid did not attain its maximum at  $(1/3, 1/3)$  but closer to the center of the domain; i.e. the refinement had led onto a different spurious branch.

NI	$\lambda$	$\ u\ $	$\dot{\lambda}$	$\dot{r}$	$\det(G_u)$	$\delta$
Level 1 solution at $r = 5$ .						
2	0.264E+00	0.500E+01	-0.255E+00	0.962E+00	0.865E 4	0.891E+00
Refine uniformly.						
Level 2 solution, no. of vertices 49.						
6	0.296E-01	0.500E+01	-0.566E-01	0.993E+00	0.530E 5	0.885E+00
Level 3 solution, no. of vertices 169.						
8	0.836E-02	0.500E+01	-0.230E-01	0.990E+00	0.626E 5	0.251E+00
Level 1 solution at $r = 4.5$ .						
2	0.431E+00	0.450E+01	-0.379E+00	0.921E+00	0.393E 4	0.824E+00
Refine adaptively.						
Level 2 solution, no. of vertices 51.						
5	0.598E-01	0.450E+01	-0.115E+00	0.983E+00	0.334E 5	0.465E+00
Level 3 solution, no. of vertices 265.						
9	0.402E-02	0.450E+01	-0.315E-01	0.862E+00	0.482E 5	0.504E-03

Table 4.2 Results of refinement on relevant branch.

One of the possible ways provided by the program to depict the results graphically is to generate a histogram. Over a square grid numbers from 0 to 9 are printed covering uniformly the range of the solution values from minimum to maximum. In Figures 4.1 and 4.2 we see which solutions were computed on the coarsest and on the finest grid for uniform and adaptive refinement.

NI	$\lambda$	$\ u\ $	$\dot{\lambda}$	$\dot{r}$	$\det(G_u)$	$\delta$
Level 1 solution at $r = 5.5$ .						
2	0.560E-01	0.550E+01	-0.772E-01	0.994E+00	-0.839E	5 0.219E+01
Refine uniformly.						
Level 2 solution, no. of vertices 49.						
9	0.328E-01	0.550E+01	-0.367E-01	0.508E+00	-0.361E	5 -0.117E+00
Level 3 solution, no. of vertices 169.						
10	0.474E-02	0.550E+01	0.112E-01	-0.990E+00	0.615E	5 0.182E+00
Refine adaptively.						
Level 2 solution, no. of vertices 41.						
4	0.968E-02	0.550E+01	-0.194E-01	0.994E+00	-0.824E	5 0.691E+00
Level 3 solution, no. of vertices 174.						
7	0.111E-03	0.550E+01	-0.386E-03	0.999E+00	0.638E	5 0.776E-01

Table 4.3 Results of refinement on spurious branch.

Norm	Relevant Solution		Spurious Solution	
	Uniform	Adaptive	Uniform	Adaptive
$H^1$	0.526	0.593	-0.955	0.448
$L^2$	1.381	1.268	-0.312	1.198
$L^\infty$	0.965	1.186	-1.099	1.183

Table 4.4 Numbers of estimated correct digits.

0000000000000000	0000000000000000	0000000000000000
0000000000000000	0000000000000000	0000000000000000
0000111111111100	0000111111100000	0000000000000000
000122222222100	000112222111000	0000111111000000
00123333332100	001123332211000	000111111110000
001234444432100	001234443321100	001122222110000
001246655432100	012345765321100	001233332211000
001367775432100	012357985432100	001244433211000
002578876432100	012358975422100	012357643211000
014689876432100	012356764321100	012359743211000
024688764332100	001234543221000	001345543211000
024666532221000	001223332211000	001233322110000
024444211110000	000111221110000	001112211100000
022221000000000	000000000000000	000001100000000
000000000000000	000000000000000	000000000000000

Figure 4.1 Histograms of the spurious solution on the coarsest grid and for uniform and adaptive refinement.

0000000000000000	0000000000000000	0000000000000000
000001111122220	0000000000000000	0000000000000000
000023333444420	000011111110000	000000111100000
000234455666420	000111222211000	000011111110000
002356677886420	001122333221100	000112222211000
013467889986410	001123454322100	000122333221100
013468899975310	001234576432100	001123454321100
013578999875310	001235797532100	001223595321100
013579998864310	001234675432100	001123564321100
014689988764310	001223454321100	001123333221000
024688776653200	001122333221100	000112222211000
024666554432000	000112222111000	000111121110000
024444333320000	000011111110000	000001111000000
022221111100000	000000000000000	000000000000000
000000000000000	000000000000000	000000000000000

Figure 4.2 Histograms of the relevant solution on the coarsest grid and for uniform and adaptive refinement.

### References

1. I. Babuska and W. C. Rheinboldt, Computational error estimates and adaptive processes for some nonlinear structural problems, *Comp. Meth. Appl. Mech. Eng.* 34 (1982), 895-937.
2. R. E. Bank, PLTMG User's Guide, Edition 4.0, Tech. Report, Dept. Math., University of California, San Diego, 1985.
3. R. E. Bank, W. M. Coughran, Jr., W. Fichtner, E. H. Grosse, D. J. Rose and R. K. Smith, Transient Simulation of Silicon Devices and Circuits, Numerical Analysis Manuscript 85-3, AT&T Bell Laboratories, Murray Hill, NJ 07974.
4. R. E. Bank and H. D. Mittelmann, Continuation and multi-grid for nonlinear elliptic systems, to appear in *Proceedings of the 2nd European Conference on Multigrid Methods*, 1985, W. Hackbusch and U. Trottenberg, eds., Springer-Verlag, Berlin.
5. R. E. Bank and A. Weiser, Some a posteriori error estimates for elliptic partial differential equations, *Math. Comp.* 44 (1985), 283-301.
6. W. J. Beyn and J. Lorenz, Spurious solutions for discrete superlinear boundary value problems, *Math. Meth. Appl. Sci.* 1 (1979), 566-571.
7. D. W. Decker and H. B. Keller, Path following near bifurcation, *Comm. Pure Appl. Math.* 34 (1981), 149-175.
8. J. Descloux, Two remarks on continuation procedures for solving some nonlinear problems, manuscript (1983)
9. H. Kuiper, On spurious numerical solutions for nonlinear eigenvalue problems, Tech. Report No. 86, Dept. Math., Arizona State University (1985).
10. H. D. Mittelmann, An efficient algorithm for bifurcation problems of variational inequalities, *Math. Comp.* 41 (1983), 473-485.
11. H. D. Mittelmann, Continuation near symmetry-breaking bifurcation points, in "Numerical methods for bifurcation problems", T. Küpper, H. D. Mittelmann and H. Weber, eds., ISNM 70, Birkhäuser-Verlag, Basel, 1984.
12. H. D. Mittelmann, Multi-level continuation techniques for nonlinear boundary value problems with parameter dependence, *Appl. Math. Comp.* (to appear).
13. H. D. Mittelmann, A pseudo-arclength continuation method for nonlinear eigenvalue problems, *SIAM J. Numer. Anal.* (to appear).
14. H. D. Mittelmann, A fast solver for nonlinear eigenvalue problems, in "Iterative solution of nonlinear systems", R. Ansorge, T. Meis and W. Törnig, eds., Springer Lecture Notes in Mathematics, vol. 953, 1982.

#### OTHER REPORTS IN THIS SERIES

Bustoz, J. and Savage, N., "Inequalities for Ultraspherical and Laguerre Polynomials," Technical Report No. 1.

Bustoz, J., "On Absolute Euler-Knopp and de la vallee Poisson Summability," Technical Report No. 2.

Busby, R.C. and Smith, H.A., "Product-Convolution Operators and Mixed-Norm Spaces," Technical Report No. 3.

Kupperman, R. H. and Smith, H. A., "The Effects of MIRV on Force Size in Mutual Deterrence," Technical Report No. 4.

Aickin, Mikel, "Linear Analysis on Contingency Tables," Technical Report No. 5.

Grace, E. E., "Refinable maps on graphs are near homeomorphisms," Technical Report No. 6.

Kuiper, H. S. and Shew, S. M., "Strong Solutions for Infinite-Dimensional Riccati Equations Arising in Transport Theory," Technical Report No. 7.

Atalla, Robert E., "Generalized Almost-Convergence vs. Matrix Summability," Technical Report No. 8.

Anderson, B. A. and Gross, K. B., "A Partial Starter construction," Technical Report No. 9.

McDonald, John N., "Approximation by quotients of rational inner functions," Technical Report No. 10.

Anderson, B. A., Gross, K. B., and Leonard, P. A., "Some Howell Designs of Prime Side," Technical Report No. 11.

Wang, Cecilia Y., "Stiffness of Harmonic Functions," Technical Report No. 12.

Wang, Alan P., "Dissipative Properties of A Nonlinear Differential Operators Equation," Technical Report No. 13.

Aickin, Mikel, "Existence of MLEs for Discrete Linear Exponential Models," Technical Report No. 14.

Wang, Cecilia Y., "A biharmonic kernel function," Technical Report No. 15.

Chung, Lung Ock, Sario, Leo, and Wang, Cecilia, "Harmonic  $L^p$  functions on the Poincare N-ball," Technical Report No. 16.

Talaga, Paul C., "Invariant Sets, Existence Theorems, and the Hukuhara-Kneser Property for Semilinear Parabolic Systems with Neumann Type boundary Condition," Technical Report No. 17.



Baumeister, Richard, "Applications of Clebsch Potentials to Variational Principles in the Theory of Physical Fields," Technical Report No. 18.

Kuiper, Hendrik J., "Invariant Sets for Nonlinear Elliptic and Parabolic Systems," Technical Report No. 19.

McMahon, Douglas, "Relativized weak mixing of uncountable order," Technical Report No. 20.

Bustoz, J. and Savage, N., "Inequalities for ultraspherical and Laguerre polynomials II," Technical Report No. 21.

Schwartz, David P., "Time-sliced Multi-tasking in a two-job Environment" (Communicated by Professor Alan Feldstein, Dept. of Mathematics, ASU, Tempe, AZ) Technical Report No. 22.

Waltman, Paul, Hubbell, Stephen P. and Hsu, Sze-Bi, "Theoretical and Experimental Investigation of Microbial Competition in Continuous Culture," Technical Report No. 23.

Anderson, B. A., "Hyperovals and Howell Designs," Technical Report No. 24.

Anderson, B. A. and Leonard, P. A., "Sequencings and Howell Designs," Technical Report No. 25.

Anderson, B. A., "A Doubling Construction for Certain Howell Designs," Technical Report No. 26.

Feldstein, A. and Schwartz, D., "A Beginner's Guide to Tape Sabatoge," Technical Report No. 27.

Leonard, Philip A. and Williams, Kenneth S., "The Spinor Genera for a Certain Form Discriminant," Technical Report No. 28.

Anderson, B. A., "Some Howell Designs of Side 6P," Technical Report No. 29.

McDonald, John N., "On Some Zero Configurations Associated with the van der Waerden Conjecture," Technical Report No. 30.

Kelly, John B., "Difference Systems, Graph Designs, and Coloring Problems," Technical Report No. 31.

Bustoz, J., "Two parameter Turan inequalities for Ultraspherical and Laguerre polynomials," Technical Report No. 32.

Ismail, Mourad E.H., "An Asymptotic Result for a Learning Model," Technical Report No. 33.

Feldstein, Alan and Neves, Kenneth W., "High Order Methods for State Dependent Delay Differential Equations With Nonsmooth Solutions," Technical Report No. 34.

Anderson, B. A. and Gross, K. B., "Asymptotic Multiplication of Howell Designs by 3," Technical Report No. 35.

Anderson, B. A., "A Note on Howell Designs," Technical Report No. 36.

Anderson, B. A. and Leonard, P. A., "Some Howell Designs of Prime Side II," Technical Report No. 37.

McDonald, J. N., "Closed Orbits of Convex Sets of Operators on the Disk Algebra," Technical Report No. 38.

Young, Dennis, "The Linear Nearest Neighbor Statistic," Technical Report No. 39.

Ismail, Mourad E.H., "The Basic Bessel Functions and Polynomials," Technical Report No. 40.

Aickin, Mikel, "DIGRAPH : A program for Testing Multiple Hypotheses," June 1980, Technical Report No. 41.

Salamon, Peter, Bjarne Andresen, Paul D. Gait, and R. Stephen Berry, "The Significance of Weinhold's Length", June 1980, Technical Report No. 42.

Salamon, Peter, and A. Nitzan, "Finite Time Optimizations of A Newton's Law Carnot Cycle," June 1980, Technical Report No. 43.

Salamon, Peter, Yehunda B. Band, and Oded Kafri, "Finite Time Thermodynamics: Optimal Expansion of A Heted Working Fluid," June 1980, Technical Report No. 44.

Salamon, Peter, Y. B. Band and O. Kafri, "Maximum Power From A Cycling Working Fluid," June 1980, Technical Report No. 45.

Salamon, Peter, Yehunda B. Band, and Oded Kafri, "Maximum Work Production from a Heated Gas in a Cylinder Equipped with a Piston," June 1980, Technical Report No. 46.

Jacobowitz, Ronald and Driscoll, Michael F., "Estimability Testability, and Parametrization in Fixed-Effects Linear Models," July 1980, Technical Report No. 47.

Dudley, David, and Wang, Alan, P., "Operator Theory on the WKB Method and Bremmer Series," July 1980, Technical Report No. 48.

Ihrig, Edwin C. and Ismail, Mourad E.H., "A q-Umbral Calculus\*", August 1980, Technical Report No. 49.

Smith, Hal, "The interaction of steady state and Hopf bifurcations in a two predator - one prey competition model," November 1980, Technical Report No. 50.

Grace, Edward E., "Local Properties in  $II_Z$  Spaces," December 1980, Technical Report No. 51.

Grace, Edward E., "Cuts points in totally non-semi-locally-connected  $II_Z$  continua," December 1980, Technical Report No. 52.

Bustoz, Joaquin and Ismail, Mourad E.H., "The associated ultraspherical polynomials and their q-analogues," February 1981, Technical Report No. 53.

Aickin, Mikel, "MLA User's Handbook," February 1981, Technical Report No. 54.

Driscoll, Michael F. and Young, Dennis L., "Some Tests for Assessing Randomness in One or More Dimensions," April 1981, Technical Report No. 55.

Buzano, Ernesto, Golubitsky, Martin, "Bifurcation Involving the Hexagonal Lattices," August 1981, Technical Report No. 56.

Anderson, B. A., "Hyperplanes and Balanced Howell Rotations," August 1981, Technical Report No. 57.

Ismail, M.E.H. and Wilson, James A., "Asymptotic and Generating Relations for the q-Jacobi and  $q_3$  Polynomials," Technical Report No. 58., August 1981.

Ismail, M.E.H. and Miller Kenneth S., "An Infinitely Divisible Distribution Involving Modified Bessel Functions", Technical Report No. 59.

Keyfitz, Barbara and Kuiper, Hendrik, "Bifurcation Resulting from Changes in Domain in a Reaction-Diffusion Equation\*," Technical Report No. 60.

Herrero, Domingo A., "Compact Perturbations of Nest Algebras, Index Obstructions and A Problem of Arveson," Technical Report No. 61.

Keyfitz, Barbara and Kuiper, Hendrik, "Bifurcation Resulting from Changes in Domain in a Reaction-Diffusion Equation\*," Technical Report No. 60.

Bustoz, J. and Ismail, Mourad E.H., "Turan Type Inequalities for Ultraspherical, Continuous  $q$ -Ultraspherical, and Hermite Polynomials," Technical Report No. 63.

Wang, Alan P., "Solutions of a Dissipative Diffusion Equation in a Hilbert Space," Technical Report No. 64, January, 1982.

Smith, Hal L., "Subharmonic Bifurcation in an S-I-R Epidemic Model", Technical Report No. 65, February 1982.

Goldstein, M. and Herrero, D., "Infinite Blaschke Products Which Almost Omit a Dense Set of Values", Technical Report No. 66, March, 1982.

Goldstein, M. and McDonald, J. N., "An extremal problem for non-negative trigonometric polynomials", Technical Report No. 67, March, 1982.

Herrero, Domingo, A., "Three Results on approximation of Hilbert Space Operators", "Economical Compact Perturbations I: Erasing Normal Eigenvalues", "Compact Perturbations of Continuous Nest Algebras", Technical Report No. 68, March, 1982.

Kuiper, Hendrik J., "Generalized Operator Riccati Equations", Technical Report No. 69, April, 1982.

Kelly, John B., "Facets of the Hamming Cone", Technical Report No. 70, May, 1982.

Herrero, D., "On Limits of Nilpotents in  $L^p$ -spaces and Interpolation of Spectra", Technical Report NO. 71, July, 1982.

Herrero, D., "Two Results on Bilateral Weighted Shifts", Technical Report No. 72, September, 1982.

Herrero, D. and McDonald, J., "On Multicyclic Operators and the Vasjuninnikol'skii Discotheca", Technical Report No. 73, September, 1982.

Herrero, D., "On Multicyclic Operators. II: Two Extensions of the Notion of Quasitriangularity", Technical Report No. 74, November 1982.

McDonald, John N., "Examples of  $RP$ -measures", Technical Report No. 75, May 1983.

Smith, Hal. L., "Normal Forms for Periodic Systems", Technical Report No. 76, July 1983.

Ismail, Mourad E.H., "On Sieved Orthogonal Polynomials", Technical Report No. 77, August 1983.

Ismail, Mourad E.H. and Askey, Richard A., "Recurrence Relations, Continued Fractions and Orthogonal Polynomials", Technical Report No. 78, September, 1983.

Mittelmann, H.D., "Continuation Near Symmetry-Breaking Bifurcation Points", Technical Report No. 79, October 1983.

Miel, George, "Perturbed Projection Methods for Split Equations of the First Kind", Technical Report No. 80, December, 1983.

Miel, George, "On the Galerkin and Collocation Methods for a Cauchy Singular Integral Equation", Technical Report No. 81, December, 1983.

McDonald, John N., "Unimodular Approximation in Function Algebras", Technical Report No. 82, March 1984.

Ringhofer, Christian A., "A Second Order Difference Scheme for Transient Semiconductor Device Simulation", Technical Report No. 83, January, 1985.

Mittelmann, Hans D., "A Free Boundary Problem and Stability for the Nonlinear Beam", Technical Report No. 84, March 1985.

Mittelmann, Hans D., "Multi-Level Continuation Techniques for Nonlinear Boundary Value Problems with Parameter Dependence", Technical Report No. 85, March 1985.

Kuiper, Hendrik J., "On Spurious Numerical Solutions for Nonlinear Eigenvalue Problems," Technical Report No. 86, July 1985.

Mittelmann, Hans D., "Continuation and Multi-grid for Nonlinear Elliptic Systems," Technical Report No. 87, September 1985.

Mittelmann, Hans D., "A Pseudo-Arclength Continuation Method for Nonlinear Eigenvalue Problems," Technical Report No. 88, October 1985.

Smith, Hal L., "Systems of ordinary differential equations which generate an order preserving flow. A survey of results.", Technical Report No. 89, December 1985.

Mittelmann, Hans D., "Multi-grid Continuation and Spurious Solutions for Nonlinear Boundary Value Problems", Technical Report No. 90, January 1986.

END

12-86

DTIC